



ORIGINAL ARTICLE

Evaluation of Antibacterial Activity of Essential Oil of *Ziziphora clinopodioides* and *Achillea wilhelmsii* on Antibiotic-resistant Strains of *Staphylococcus aureus*

Abolfazl Jafari-Sales<sup>1,2\*</sup>, Ahmadreza Shahniani<sup>2</sup>, Reza Fathi<sup>3</sup>, Parviz Malekzadeh<sup>4</sup>, Haedeh Mobaiyen<sup>5</sup>, Farnaz Rasi-Bonab<sup>6</sup>

<sup>1</sup>Young Researchers and Elite Club, Ahar Branch, Islamic Azad University, Ahar, Iran

<sup>2</sup>Department of Microbiology, Kazeroon branch, Islamic Azad University, Kazeroon, Iran

<sup>3</sup>Department of Physiology, School of Medicine, Marmara University, Istanbul, Turkey

<sup>4</sup>Department of Biology, Faculty of Science, Qom University, Qom, Iran

<sup>5</sup>Department of Microbiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran

<sup>6</sup>Young Researchers and Elite Club, Marand Branch, Islamic Azad University, Marand, Iran

**Corresponding Author:** Abolfazl Jafari Sales, E-mail: a.jafari\_1392@yahoo.com

ARTICLE INFO

Article history

Received: Mar 29, 2017

Accepted: Apr 17, 2017

Published: May 23, 2017

Volume: 2

Issue: 2

Conflicts of interest: None

Funding: None

Key words

Antibacterial Effects,

Essential Oil,

*Staphylococcus aureus*

ABSTRACT

**Background:** The use of drugs to treat diseases led to the emergence of resistant strains of bacteria. Bacterial resistance to antibiotics is the most common problems in medical science. This study was aimed to evaluate the antimicrobial effects of essential oil of *Ziziphora clinopodioides* and *Achillea wilhelmsii* on against resistant clinical strains of *Staphylococcus aureus*. **Methods:** After collecting plants and validate its scientific name by botanists of Agricultural Organization and after drying in the shade, essential oil of *Z. clinopodioides* and *A. wilhelmsii*, extracted with steam distillation method by Clevenger and antimicrobial effects of essential oil by well diffusion on bacteria mentioned above were interpreted. The amount of essential oil were injected to gas chromatography linked to mass spectrometry (GC/MS), and the amount and type compounds of the essential oils were identified. **Results:** The results showed that the extracted essence essential oil from the *Z. clinopodioides* and *A. wilhelmsii* has a bactericidal effect. In the obtained results by GC/MS chromatography, in *Z. clinopodioides* and *A. wilhelmsii* essential oils, 22 and 18 compounds were identified, respectively. **Conclusion:** Our results showed *Z. clinopodioides* and *A. wilhelmsii* have considered in herbals groups with antibacterial properties and after evaluating their effects *in-vivo* condition and to identify the active ingredients, as an alternative to synthetic drugs that commonly used to treat infections are used.

INTRODUCTION

One of the most important health challenges is the treatment of infectious diseases due to their much higher prevalence and spread. Since the discovery of penicillin in the late 40's, and widely use of it, new antibiotics have been continuously presented to treat infections. The result was the increased use of clinical or synthetic antibiotics to heal clinical infectious disease. The indiscriminate use of antimicrobial drugs caused the increased drug resistance against different antibiotics in most bacteria (1-6). It has been one of the reasons of the increasing use of plants as natural, safe, accessible and affordable materials in treating bacterial infections in comparison with synthetic antibiotics. Also, the consumption of herbal medicines is more acceptable than chemical drugs (7-9). Moreover, that is why a new wave of expensive global studies has raised, and the antibacterial effects of dif-

ferent plants have been introduced (10). Therefore, two plant species were considered in the current study.

*Ziziphora clinopodioides* belongs to *Ziziphora* genus and Lamiaceae family with the Persian name of *kakuti-e kuhi* is a perennial plant. In Iran, several hundred species in 49 genera of the Lamiaceae family are scattered. The geographical distribution of *Z. clinopodioides* in the world is as following: eastern Balkan peninsular, southeast of Asia and from central Asia to Pamir-Alay and Himalaya mountains (Iran, Iraq, central and east parts of Turkey) and Africa (11). Chitsaz et al. (2005) examined the compounds and antimicrobial potential of *Z. clinopodioides* essential oil on several bacterial species. The essential oil of the plant prevents the growth of all studied gram-negative and gram-positive organisms and had the most effect on *Salmonella typhimurium* (12). Salehi et al. (2006) studied the antibacterial activity and chemical compounds of *Z. clinopodioides* essential oil.

Thirty-two components constituting 97.1 % of the essential oil were identified (13). *Achillea wilhelmsii* C.Koch is one of the most famous medicinal plants that about 85 species of it have been identified that 7 of them are exclusively found in Iran. Native *Achillea* species has a relatively wide distribution in different parts of the country and are commonly grown in fields, roadsides and mountainous areas (14). New flowers of *Achillea* are rich in chemicals and used to treat respiratory problems as anti-allergic, anti-congestion and mucus. Essential oil of them contains chamazulene, cineol and borneol which have shown anti-inflammatory, antispasmodic, anti-dandruff properties, hair growth stimulation, and topical healing of skin (14-15).

Aljancic et al. (1999) in their study showed the significant inhibitory effect of *A. wilhelmsii* on *Candida albicans* and *Bacillus subtilis*. According to them, in addition to the inhibitory effect of two above microbes, the flavonoids of *A. wilhelmsii* essential oil also prevent the growth of *Aspergillus Niger* effectively (16).

In the study by Sokmen et al. (2003), the antimicrobial effect of the essential oil, aqueous and methanol extracts of *A. sintenisii* on 12 bacteria and two yeast species were examined comparatively (17). The current study intends to investigate the chemical compounds and anti-bacterial effects of *Z. clinopodioides* and *A. wilhelmsii* on *Staphylococcus aureus* resistant to the antibiotic.

## METHODS

### 1. Plant sampling and investigation of its anti-microbial properties

Plant samples were collected from the nature areas of Eyshabad village, Marand city, and East-Azerbaijan province in two stages in late May to early June. The samples were cleaned after collection and transmission, and also, they were dried in a large and proper space under the conditions away from direct sunlight. After they had been dried completely, their aerial parts (stems and leaves) were separated from the roots and prepared to be grinded. Then, the essential oil of their aerial parts was extracted for 5 hours by a Clevenger-type apparatus. After washing with sodium sulfate and dissolving in n-hexane, their essential oil was used to investigate anti-bacterial properties. The bacteria employed in this study was *S. aureus* isolated from patients. Initially, a suspension equivalent to standard McFarland 0.5 turbidity was prepared from each bacterium. After spreading the bacterial suspension on Müller-Hinton agar media, wells of 5 mm in diameter were immediately made. About 50 µL and 100 µL of essential oils were added aseptically in wells. The same amount of n-hexane solvent (as con-

trol) was added to plates and then were incubated for 24 hours at 37°C (18-19).

### 2. Bacterial isolates and their resistance to antibiotics

Thirty samples used in this study were selected from the clinical samples of Marand Razi hospital and cultured on mannitol salt agar and blood agar media. Isolated pure colonies of bacteria on culture media were detected regarding type through catalase tests and other biochemical tests. Finally, *S. aureus* isolates were recognized by using tube coagulase test, LAM test, and agglutination investigation. The antibiotics sensitivity/resistance profiles (antibiogram) of the purified isolate were determined by the methodology of Kirby-Bauer's on Mueller-Hinton Agar medium. We used antibiotics include trimethoprim/sulfamethoxazole (SXT10 + 10 µg), ampicillin (AM 10µg), ceftazidime (SAZ 30µg), ampicillin (SXT10+10 µg), ceftazidime (CF 5 µg), amikacin (AN 30 µg), penicillin (10 µg P), ceftriaxone (30 µg CRO), erythromycin (15 µg E) and tetracycline (TE 30 µg) made by Padtan Teb Co. The diameter of inhibition zone was measured by comparison to the standard isolate after 24 hours of incubation at 37°C, and the sensitivity and resistance of isolates were determined and compared with standard table.

### 3. Essential oil analysis by gas chromatography-mass spectrometry (GC/MS)

The essential oil sample was injected in GC/MS (type: Agilent GC (USA)) after finding thermal planning system. The compounds were identified by the database of the device and comparing their mass spectra. SPSS V.19 software performed variance analysis of data, and Duncan's multiple range test at the confidence level of 99% was used for the mean.

## RESULTS

### 1. Bacterial responses to antibiotics

In-vitro antibacterial activities of some antibiotics against obtained isolates were evaluated. The results revealed that in the presence of antibiotics, each bacterial isolate responded somewhat similarly, however, there were differences among the isolates in terms of antimicrobial resistance or antimicrobial sensitivity. Some differences were also observed in inhibition mechanisms of antibiotics, as the antibacterial effect of penicillin and erythromycin was much less than other antibacterial drugs (Table 1).

### 2. Antimicrobial activity of essential oils

Because of various geographical and climatic conditions in Iran, there is a rich and varied source of plant species in it.

Some of these herbal plants have medicinal properties

**Table 1.** Antibiotic susceptibility pattern of *S. aureus* isolates (%)

	CAZ	AM	TE	STX	CF	AN	CRO	E	P
R	33.3	70	23.3	76.7	80	23.3	36.7	83.3	83.3
S	50	10	26.7	20	10	76.7	16.7	10	6.7
I	16.7	20	50	3.3	10	0	46.6	6.7	10

R: Resistant; S: Sensitive; I: Intermediate

such as antibacterial activity. In our study, the antibacterial effects of essential oils were as follows: essential oils of *Z. clinopodioides* and *A. wilhelmsii* could inhibit *S. aureus* growth.

The antibacterial effect of *Z. clinopodioides* was more than the one of *A. wilhelmsii*. The results of the antimicrobial activity of *Z. clinopodioides* and *A. wilhelmsii* are represented in Table 2.

The n-hexane solvent was used as control sample that showed no inhibitory effect. Given the observed clear relationship between the plant derivate (essential oils) and their antimicrobial activity, it can be said that they can be used as new biologically active substances against some virulent infectious agents.

**Table 2.** Antimicrobial effect of essential oils of *Z. clinopodioides* and *A. wilhelmsii* on *S. aureus* resistant to antibiotics (mean±SD)

Plant	Diameters of the inhibition zones (mm)		
	<i>S. aureus</i>		
	50 µl	100 µl	N-Hexane control
<i>A. wilhelmsii</i>	0.47±10.66	0.16±13.5	0
<i>Z. clinopodioides</i>	0.81±11	0.18±14.6	0

### 3. Chemical compounds obtained from the essential oils of *Z. clinopodioides* and *A. wilhelmsii*

Chemical compounds obtained from the essential oils of the aerial part of *Z. clinopodioides* and *A. wilhelmsii* were investigated *in-vitro*. The essential oils were extracted by steam distillation with Clevenger and injected to GC/MS after purification (removing from the plant sweat), and the amount and type of the essential oil were determined. Totally, 22 compounds of *Z. clinopodioides* essential oil, including 95.4% of compounds, and 18 compounds of *A. wilhelmsii* essential oils, including 91.31% of compounds, were identified. Table 3 shows the ingredients of each essential oil.

## DISCUSSION

According to increased bacterial resistance to a variety of antibiotics, it has been tried to access and use the compounds of plants in the treatment of various diseases. From thousands of years ago, plants have played a critical role in maintaining health and improving the life quality of human beings. Medicinal herbs have useful properties that their antibacterial, anti-parasitic, antifungal and antioxidant properties can be noted (20).

Chalibian et al. (2003) have proved that *A. wilhelmsii* has an antimicrobial effect on *S. aureus* that the results of this study are consistent with it (21).

**Table 3.** Chemical compounds obtained from the essential oils of *Z. clinopodioides* and *A. wilhelmsii*

<i>A. wilhelmsii</i>			<i>Z. clinopodioides</i>		
#	Compounds	Percent	#	Compounds	Percent
1	alpha-Pinene	18.82	1	Alpha-thujene	0.1
2	beta-Pinene	11.2	2	alpha-Pinene	3.8
3	Caryophyllene oxide	9.21	3	Camphene	0.4
4	Alpha-thujene	8.62	4	Sabinene	4.3
5	Valencene	7.92	5	beta-Pinene	4.3
6	Chamazulene	6.31	6	Myrcene	1.8
7	ρ-Cymene	5.82	7	1,8-cineole	19.7
8	Aromadendren	5.45	8	beta oemene	0.3
9	Camphor	4.74	9	Terpinene	0.9
10	delta-cadinene	4.12	10	Isomenthone	1.2
11	gamma-terpinene	1.94	11	Neomentho	6.9
12	Bornyl acetate	1.51	12	Neoisomenthol	0.4
13	Trans carveol	1.41	13	α-Terpineol	3.7
14	gamma- cadinene	1.32	14	Pulegone	25.9
15	Terpinolen	1.03	15	Piperitone	2.7
16	1-alpha-terpineol	0.9	16	Bornyl acetate	2.6
17	Cis carveol	0.35	17	Methyl acetate	1.1
18	Eugenol	0.32	18	Piperitenone	4.4
19	Carvacrol	0.32	19	Borneol	1.5
			20	E-Caryophyllene	3.2
			21	Spathulenol	3.6
			22	Terpinene	2.6
Total		91.31	Total		95.4

Ghaderi et al. (2012) shown that *A. wilhelmsii* had inhibitory effect on *Escherichia coli*, *Pseudomonas aeruginosa*, and *Bacillus cereus* and used GC/MS to separate and identify the compounds of essential oil that the results of the chemical compounds of the current study are consistent with the results of their study (22). In the study by Aljancic et al., it has been found that *A. wilhelmsii* prevents the growth of *Bacillus subtilis*, *C. albicans* and *A. niger* in *in-vivo* conditions. They have identified the flavonoids extracted from the essential oils as a factor of antibacterial effects (23).

Kermanshah et al. have studied the antibacterial effect of the hydroalcoholic extracts of common sage (*Salvia officinalis*) and *A. wilhelmsii* on cariogenic microorganisms and concluded that both essential oils have an inhibitory effect on the growth of *Streptococcus mutans*, *Lactobacillus rhamnosus* and *Actinomyces viscosus* (24).

Khanaki et al. (2010) have identified the chemical compounds of *Z. clinopodioides* by GC/MS. The results of the current study are consistent with their results (25). Ousalah et al. (2006) have identified the main compounds of *Z. clinopodioides* essential oil. Pulegone is a ketone and of Monoterpene. Terpenes can damage the cell membrane and penetrate the lipid structure of the cellular wall of bacteria that this leads to protein denaturation and the collapse of the cell structure and leakage of cytoplasm and finally cell death (26). Previous studies indicated that the extracts of *Z. persica* and *Z. clinopodioides* could prevent the growth of broad range of gram-positive and gram-negative bacteria (27,28).

## CONCLUSION

In general, *in-vitro* investigations showed that of the essential oils extracted from *Z. clinopodioides* and *A. wilhelmsii* have antimicrobial activity against the *S. aureus* resistant to antibiotics that would be considered as an alternative option in producing herbal medicines against the *Staphylococcus aureus* bacterium with minimal side effects. Since that they are non-toxic and natural components after performing further investigation on experimental animals.

## ACKNOWLEDGMENTS

We would like to thank the Department of Microbiology, Islamic Azad University Marand and Tabriz for their laboratory equipment and technical support.

## AUTHORS CONTRIBUTION

The authors had equal contributions.

## REFERENCES

1. Nimmo GR, Cooms GW, Pearson JC, et al. Methicillin resistant *Staphylococcus aureus* in the Australian community; an evolving epidemic. Med J Aust. 2006;184(8):374-5.
2. Orrt FA, Land M. Metecillin resistant *Staphylococcus aureus* prevalence: Current susceptibility patterns in Trinidad. BMC Infect Dis. 2006;6:83.
3. Skaltsa H, Lazari DM, Chinou IB, Loukis AE. Composition and antibacterial activity of the essential oils of *Stachys candida* and *S. chrysantha* from Southern Greece. Planta Med. 1999;65(3):255-6.
4. Skaltsa HD, Demetoz C, Lazari, Sokovic M. Essential oil analysis and antimicrobial activity of eight *Stachys* species from Greece. Phytochemistry. 2003;64(3):743-52.
5. Digrak M, Alma MH, Ilcim A. Antibacterial and antifungal activities of Turkish medicinal plants. Pharm. Biol. 2001;39(5):346-50.
6. Weinstine RA. Controlling antimicrobial resistance in hospitals: Infection control and use of antibiotics. Emerg Infect Dis 2001;7:188-92.
7. Mosaddegh M. and Naghibi FI. Traditional Medicine: Past & Present. Traditional Medicine & Materia medica. Vol. 1. Tehran, Iran; Published TMRC; 2002:p:2-20. (Persian).
8. WHO Traditional Medicine Strategy 2002-2005, Geneva. 2002;1-3:43-47.
9. The promotion & development of traditional medicine- Report of a WHO meeting” WHO Report series, No.622, Switzerland. 1978;8-13: 36-9.
10. Marjorie MC. Plant Products as Antimicrobial Agents. Clin Microb Rev. 1999;12: 564-82.
11. Baser KHC, Sezik E, Tumen G. Composition of The Essential Oil of *Ziziphora clinopodioides* Lam. Journal of Essential Oil Research. 1991;3(4):237-9.
12. Chitsaz M, Barton M, Bazargan M. Kamallnejad M. Essential oil composition and antibacterial effects of *ziziphora clinopodioides*. Journal of Shahed university. 2005;68,15-23.
13. Salehi P, Mirjalili M.H, Hadian J, Nejad Ebrahimi S, Yousef Zadi M. Antibacterial activity and composition of the essential oil of *Ziziphora clinopodioides subsp bungeana* (JUZ.). Rech. F. from Iran. Z. 2006;61:677-80.
14. Bafrani HH, Parsa Y, Yadollah-Damavandi S, Jangholi E, Ashkani-Esfahani S, Gharehbeglou M. Biochemical and Pathological Study of Hydroalcoholic Extract of *Achillea millefolium* L. on Ethylene Glycol-Induced Nephrolithiasis in Laboratory Rats. N Am J Med Sci. 2014;6(12):638-42.
15. Perry I. Medical plants of east and southeast Asia. Cambridge: Mit Press; 1980.
16. Aljancic I, Vajs V, Menkovic N, et al. Flavones and sesquiterpene lactones from *Achillea atrata* subsp multifida: Antimicrobial activity. J Natur Products. 1999;62(6):909-11.
17. Sokmen A, Sokmen M, Daferera D, et al. The in vitro antioxidant and antimicrobial activities of the essential oil and methanol extracts of *Achillea biebersteini* Afan. (Asteraceae). Phytother Res 2004;18(6):451-6.
18. Negahban, M., Moharrampour, S. and Sefidkon, F. Fumigant toxicity of essential oil from *Artemisia sieberi* Besser against three stored-product insects. Journal of Stored Product Research, 2007;43:123-8.
19. Jaymand K, Rezaee M. Essential oil, Distillations Apparatuses, test Method of essential oil and Retention Indices in Essential oil Analysis. Iranian Society of Medicines

- nal Plants. Tehran. 2007; pp:106 - 8.
20. Mirmalek SA, Azizi MA, Jangholi E, Yadollah-Damavandi S, Javidi MA, Parsa Y, et al. Cytotoxic and apoptogenic effect of hypericin, the bioactive component of *Hypericum perforatum* on the MCF-7 human breast cancer cell line. *Cancer Cell Int.* 2015;16:3.
  21. Chalabian F, Norouzi Arasi H, Moosavi S. A study of growth inhibitory effect of essential oils of seven species from different families on some kinds of microbes. *Journal of Medicinal Plants*, 2003(Issue 7);7(2):37-42. (In Persian).
  22. Ghaderi S, Falahati-Hosseini A, Sarailoo MH, Ghanbari V. Investigation of the components and antibacterial effects of three plant's essential oil *Coriandrum sativum*, *Achillea millefolium*, *Anethum graveolens* in vitro. *Journal of Shahrekord University of Medical Sciences.* 2012;14(5):74-82. (In Persian).
  23. Aljancic I, Vajs V, Menkovic N, et al. Flavones and sesquiterpene lactones from *Achillea atrata* subsp. *multifida*: Antimicrobial activity. *J Natur Products.* 1999;62(6): 909-11.
  24. Kermanshah H, Hashemi-Kamangar S, Arami S. Antibacterial activity of hydroalcoholic extract of *Salvia officinalis* and *Achilles millefolium* against cariogenic microorganisms: An In-vitro investigation. *Iran J Dentist* 2009;21(3): 215-20. (In Persian).
  25. Bakhshi Khaniki G, Sefidkon F, Dehghan Z. The effects of some ecological factors on essential oil yield and composition of *Ziziphora clinopodioides* Lam. *Journal of herbal drugs, Iran.* 2010;1;9-6. (In Persian).
  26. Oussalah M, Caillet S, Lacroix M. Mechanism of action of *Spanish oregano*, *Chinese cinnamon*, and *savory* oils against cell membrane and walls of *Escherichia coli O157:H7* and *Listeria monocytogenes*. *Journal of Food Protection.* 2006;69(5):1046-55.
  27. Ozturk S, Ercisli S. Antibacterial activity and chemical constitutions of *Ziziphora clinopodioides*. *Food Control.* 2007;18(5):535-40.
  28. Ozturk S, Ercisli S. The chemical composition of essential oil and in vitro antibacterial activities of essential oil and methanol extract of *Ziziphora persica bunge*. *Journal of Ethnopharmacology.* 2006;106(3):372-6.